

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Auctores Latini Pedemontani. Un'antologia degli scrittori in lingua latina in Piemonte fra Ottocento e Novecento**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1737632> since 2020-04-28T22:23:22Z

*Publisher:*

Dell'Orso

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in:

*Journal of Asian Earth Sciences* (2011), 42, 4: p. 607–17, doi:10.1016/j.jseaes.2010.10.016.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform*]

Manuscript Number: JAES-D-10-00055

Title: Ultrahigh-pressure metamorphism and multistage exhumation of eclogite from the Luotian dome, North Dabie Complex Zone (central China): Evidence from mineral inclusions and decompression texture

Article Type: SI: HP-UHP metamorphic rocks

Keywords: Eclogite; Coesite; Ultrahigh-pressure metamorphism; Multistage exhumation; North Dabie Complex Zone

Corresponding Author: Professor Yican Liu, Ph.D.

Corresponding Author's Institution:

First Author: Yican Liu, Ph.D.

Order of Authors: Yican Liu, Ph.D.; Xiaofeng Gu; Franco Rolfo; Zhenyu Chen

**Abstract:** The studied eclogite is located in the Luotian dome in the southwestern part of the North Dabie Complex Zone (NDZ), central China, and is a portion of deeply subducted mafic lower continental crust of the South China Block. Petrologic analysis suggests that the eclogite underwent ultrahigh-pressure (UHP) and high-pressure eclogite-facies metamorphism, and subsequent HP granulite-facies overprinting and amphibolite-facies retrogression during continental subduction and exhumation. As a result, multiple decompression textures were produced, suggesting a multistage exhumation history from eclogite-, to granulite-, to amphibolite-facies conditions. A striking feature of this eclogite is the widespread exsolution of different phases: oriented needles of rutile + clinopyroxene + amphibole + apatite in garnet; quartz + amphibole + hyperthsenite + sodic plagioclase in clinopyroxene; pyrrhotite in apatite. Most importantly, we provide for the first time conclusive evidence of quartz pseudomorphs after coesite in garnet and relic coesite (confirmed by in situ Raman spectroscopy) in zircon. In addition, there are two groups of apatite: one is fluor (F)-apatite and occurs as inclusion in garnet or in matrix, which is rich in F ( $\geq 3$  wt %) without any exsolutions and formed at UHP metamorphic conditions; the second is relatively poor in F ( $< 1$  wt %) associated with the oriented pyrrhotite exsolution and formed during decompression. Microtextural and petrologic analysis suggest that the eclogites from southwestern segment of the NDZ are similar to eclogites from the northern segment of the NDZ (e.g., Huangweihe and Baizhangyan) and suffered UHP metamorphism with pressure peak  $\geq 5$ -7 GPa. These results, combined with published geochronological data, imply that the NDZ wholly experienced Triassic UHP metamorphism as a coherent unit.

1        **Ultrahigh-pressure metamorphism and multistage exhumation of**  
2        **eclogite from the Luotian dome, North Dabie Complex Zone (central**  
3        **China): Evidence from mineral inclusions and decompression texture**

4  
5                Yi-Can Liu <sup>a,\*</sup>, Xiao-Feng Gu <sup>a</sup>, F. Rolfo <sup>b</sup> and Zhen-Yu Chen <sup>c</sup>

6  
7        a. CAS Key Laboratory of Crust-Mantle Materials and Environments, School of Earth and Space  
8        Sciences, University of Science and Technology of China, Hefei 230026, China

9        b. Department of Mineralogical and Petrological Sciences, University of Torino, Via Valperga  
10       Caluso 35, I-10125 Torino, Italy

11       c. Institute of Mineral Resources, Chinese Academy of Geological Sciences, Beijing 100037,  
12       China

13  
14  
15  
16  
17  
18  
19       \*Corresponding author. Tel.: +86 551 3600367.

20       *E-mail address:* [liuyc@ustc.edu.cn](mailto:liuyc@ustc.edu.cn) (Yi-Can Liu)

21

## ABSTRACT

The studied eclogite is located in the Luotian dome in the southwestern part of the North Dabie Complex Zone (NDZ), central China, and is a portion of deeply subducted mafic lower continental crust of the South China Block. Petrologic analysis suggests that the eclogite underwent ultrahigh-pressure (UHP) and high-pressure eclogite-facies metamorphism, and subsequent HP granulite-facies overprinting and amphibolite-facies retrogression during continental subduction and exhumation. As a result, multiple decompression textures were produced, suggesting a multistage exhumation history from eclogite-, to granulite-, to amphibolite-facies conditions. A striking feature of this eclogite is the widespread exsolution of different phases: oriented needles of rutile + clinopyroxene + amphibole + apatite in garnet; quartz + amphibole + hypersthene + sodic plagioclase in clinopyroxene; pyrrhotite in apatite. Most importantly, we provide for the first time conclusive evidence of quartz pseudomorphs after coesite in garnet and relic coesite (confirmed by in situ Raman spectroscopy) in zircon. In addition, there are two groups of apatite: one is fluor (F)-apatite and occurs as inclusion in garnet or in matrix, which is rich in F (~3 wt %) without any exsolutions and formed at UHP metamorphic conditions; the second is relatively poor in F (< 1 wt %) associated with the oriented pyrrhotite exsolution and formed during decompression. Microtextural and petrologic analysis suggest that the eclogites from southwestern segment of the NDZ are similar to eclogites from the northern segment of the NDZ (e.g., Huangweihe and Baizhangyan) and suffered UHP metamorphism with pressure peak > 5-7 GPa. These results, combined with published

44 geochronological data, imply that the NDZ wholly experienced Triassic UHP  
45 metamorphism as a coherent unit.

46

47 **Key words:** Eclogite; Coesite; Ultrahigh-pressure metamorphism; Multistage  
48 exhumation; North Dabie Complex Zone

## 1. Introduction

The oriented exsolution or lamellae is a common feature of minerals that derive from deep-seated sources such as kimberlites, diamondiferous rocks, as well as eclogites and related rocks from high-pressure (HP) to ultrahigh-pressure (UHP) metamorphic belts, and is interpreted as the result of cooling and decompression (e.g., Smith, 1984; Gayk et al., 1995; Zhang et al., 1995, 2002a, 2005; Katayama et al., 2000a; Tsai and Liou, 2000; Ye et al., 2000a; Dobrzhinetskaya et al. 2002; Song et al., 2003, 2005; Malaspina et al., 2006; Nakano et al., 2007 and references therein). This peculiar microtexture is generally used to infer the primary conditions under which eclogites and related metamorphic rocks were formed. Eclogites are considered to be the most important witnesses of continental subduction, collision and subsequent exhumation processes and thus provide important information on the geodynamics of orogens. The survival of coesite and other index minerals in UHP rocks has important implications for the exhumation of subducted crustal rocks and is most commonly attributed to rapid exhumation, continuous cooling during uplift, and inclusion in strong phases that can sustain a high internal over-pressure during decompression (Mosenfelder et al., 2005). However, peak metamorphic conditions and the pressure/temperature (PT) evolution of UHP rocks involved in multistage metamorphism are not always straightforward because their mineral assemblages have been re-equilibrated to various degrees during retrogression and decompression (e.g., Faryad, 2009).

The finding of coesite and micro-diamond in eclogites from the Dabie orogen in

central China (Okay et al., 1989; Wang et al., 1989; Xu et al., 1992) led this area to be one of the most important targets for studying UHP metamorphism, documenting continental subduction to mantle depths. A great number of studies have contributed to the understanding of the geodynamics of subduction and exhumation resulting in the Dabie–Sulu orogenic belt (see Liu and Li, 2008 for a summary). This orogenic belt contains the largest exposure of UHP rocks in the world, which formed by Triassic continental collision between the South China and North China Blocks (e.g., Li et al., 1993, 2000; Chavagnac and Jahn, 1996; Hacker et al., 1998; Ayers et al., 2002; Liu et al., 2005, 2007a, 2007b).

The Dabie UHP metamorphic belt in central China consists of three eclogite-bearing UHP crustal slices from north to south: the North Dabie complex zone (NDZ), the Central Dabie UHP metamorphic zone (CDZ), and the South Dabie low-T eclogite zone (SDZ), which differ by rock association, protolith nature and metamorphic evolution (see Liu and Li, 2008, in detail). Moreover, Pb isotopic mapping on the Dabie UHP belt has revealed a clear detachment between deeply subducted upper continental crust (the CDZ) and felsic lower crust (the NDZ) (Zhang et al., 2002a; Li et al., 2003). This duality is also supported by their different termination ages of peak UHP metamorphism, which probably result from multistage detachment within deeply subducted crust at different depths and multi-slice successive exhumation of the UHP rocks during continental collision (Liu et al., 2007b; Liu and Li, 2008). However, because of the lack of critical UHP phases such as coesite or microdiamond in the southwestern part of the NDZ, whether or not the



NDZ as a whole underwent deep subduction and subsequent UHP metamorphism is still a controversial issue (Zhao et al., 2008; Zhang et al., 2009).

In order to clarify : (1) peak metamorphic conditions of the eclogite from the Luotian dome in the southwestern part of the NDZ; (2) if the NDZ wholly underwent UHP metamorphism, a petrologic study on the eclogite from the Luotian dome in the NDZ was carried out. This paper focuses on multiple decompression texture and UHP metamorphic evidence of these eclogites. The results not only provide direct evidence on UHP metamorphism and multistage breakdown processes on the eclogite in the area, but also provide new constraints on exhumation mechanism of the UHP rocks in the Dabie orogen.

## **2. Geologic background and sample**

The Dabie orogen is located in the intermediate segment of the Qinling-Dabie-Sulu belt formed by the collision of the North China Block and South China Block (SCB) in the Triassic. It comprises several fault-bounded rock units with varying metamorphic grades and is generally subdivided into five major lithotectonic units from north to south (e.g., Okay et al., 1993; Xu et al., 2003, 2005; Li et al., 2004; Liu et al., 2005, 2007a, 2007b; Liu and Li, 2008): (1) the Beihuaiyang zone (BZ); (2) the North Dabie complex zone (NDZ); (3) the Central Dabie UHP metamorphic zone (CDZ); (4) the South Dabie low-T eclogite zone (SDZ); and (5) the Susong complex zone (SZ). These five zones are respectively separated by the Xiaotian-Mozitan fault (XMF), Wuhe-Shuihou fault (WSF), Hualiangting-Mituo fault (HMF) and

Taihu-Shanlong fault (TSF) (Fig. 1). Zone (1) is a low-grade composite unit comprising the Foziling (or Xinyang) Group and the Luzhenguan (or Guishan) complex, whereas Zones (2), (3), (4) and (5) belong to the subducted SCB.

UHP metamorphic rocks, including coesite-bearing eclogite, UHP gneiss, whiteschist, quartz jadeitite and marble with eclogite nodules, are observed in the CDZ and SDZ (e.g., Xu et al., 1992; Okay, 1993; Okay et al., 1993; Rolfo et al., 2000; Li et al., 2004). The occurrence of diamond and coesite in the UHP rocks from the CDZ indicates the UHP metamorphism occurred at 700–850 °C and >2.8 GPa (e.g., Okay et al., 1989; Wang et al., 1989; Xu et al., 1992; Okay, 1993; Rolfo et al., 2004), whereas the peak P-T conditions on the eclogites in the SDZ were estimated at 670 °C and 3.3 GPa (Li et al., 2004). Both the CDZ and SDZ units experienced UHP eclogite-facies, and subsequent HP eclogite- and amphibolite-facies retrograde metamorphism (e.g., Xu et al., 1992; Okay, 1993; Rolfo et al., 2004; Li et al., 2004).

The NDZ consists predominantly of banded tonalitic and granitic gneiss and post-collisional intrusions with subordinated meta-peridotite (including dunite, harzburgite and lherzolite), garnet pyroxenite, garnet-bearing amphibolite, granulite and eclogite. Although no coesite has not yet been discovered from the eclogite or gneiss in the NDZ, the occurrence of micro-diamond from eclogite and gneiss is a key evidence for UHP metamorphism (Xu et al., 2003, 2005; Liu et al., 2007b). In addition, quartz rods in clinopyroxene are widespread in the eclogite, and these exsolutions are commonly regarded as evidence of prior UHP metamorphism (e.g., Smith, 1984; Liou et al., 1998; Tsai and Liou, 2000; Liu et al., 2005, 2007a; Zhang et

al., 2005). Therefore, also the eclogites from the NDZ underwent the UHP metamorphism at  $P > 3.5\text{--}4.0$  GPa (Xu et al., 2003, 2005; Malaspina et al., 2006). As concerning geochronology, Triassic zircon U-Pb (Liu et al., 2000) and Sm-Nd (Liu et al., 2005) ages of the eclogites from the NDZ suggest that they formed by subduction of the SCB in the Triassic, similar petrogenesis to those from the CDZ and SDZ. The Triassic metamorphic ages (Liu et al., 2000, 2007b; Bryant et al., 2004; Xie et al., 2004) and the occurrence of micro-diamonds in zircons (Liu et al., 2007b) from the banded gneisses in the NDZ suggest that also the gneisses surrounding the eclogites were involved in the continental deep subduction of the SCB. Following the UHP and HP eclogite-facies metamorphism, however, the eclogites from the NDZ were first subjected to granulite-facies overprinting, and later to amphibolite-facies retrogression (e.g., Liu et al., 2000, 2005; Xu et al., 2000). This peculiar P-T path suggests a different post-peak metamorphic evolution for different tectonic units (e.g., NDZ and CDZ) of the Dabie UHP belt. Therefore, although both the CDZ and NDZ units experienced UHP metamorphism, they had different exhumation histories, suggesting that CDZ and NDZ are two UHP slices decoupled after subduction (Xu et al., 2005; Liu et al., 2005, 2007a, 2007b; Liu and Li, 2008). As concerning the metamorphic history prior to the peak, Pb isotope investigations show that the UHP rocks from the CDZ are characterized by high radiogenic Pb while banded gneiss from the NDZ are characterized by low radiogenic Pb (Zhang et al., 2002a; Li et al., 2003); this suggests that the UHP rocks of the CDZ were derived from subducted upper crust, while the UHP rocks of the NDZ were derived from subducted felsic lower continental crust

with minor mafic boudins or lenses (Li et al., 2003; Liu et al., 2007a, 2007b).

A peculiar geologic feature in the southwestern segment of the NDZ is the Luotian dome, which is a deeply eroded area with abundant felsic and mafic granulites around the Luotian county (Fig. 1). In the Luotian dome, unusual eclogites were found as lenses in garnet-bearing banded tonalitic gneisses (Liu et al., 2007a). These eclogites preserve early granulite-facies mineral relics and have been overprinted by regionally extensive HP granulite-facies metamorphism, followed by penetrative amphibolite-facies retrogression during exhumation. The eclogite-facies mineral assemblage is garnet and relict omphacite, with rutile, quartz, allanite and fluor-apatite as common constituents. While micro-diamonds were found in the northeastern part of the NDZ, no index UHP minerals such as coesite and micro-diamond were identified so far in southwestern part of the NDZ, i.e. in the Luotian dome area. Because of the lack of index UHP phases, it has been always difficult to determine accurately the peak conditions under which the eclogites were formed, thus feeding a number of arguments on the tectonic affinity and evolution of the NDZ.

The eclogites from the Luotian dome occur as bands, lenses and blocks, up to 3m thick (Fig. 2), and are often strongly retrogressed to (garnet) amphibolite. The studied samples were collected from Luotian (sample 03LT1-1), Jinjiapu (samples 03LT8-1, 06LT3-2 and 06LT4-2) and Shiqiaopu (samples 07LT6-1 and 09LT2-1), respectively (Fig. 1). Details of the petrography and mineral chemistry of the studied samples were given in a separate paper (Liu et al., 2007a) and are only summarized here. However,

samples were chosen because of their relative abundance of mineral inclusions in zircon and garnet, and oriented mineral exsolutions in garnet, clinopyroxene and apatite. Sample 03LT1-1 is a strongly retrogressed eclogite, which is mainly composed of garnet, rutile, hornblende and plagioclase with minor quartz, diopside, hypersthene and ilmenite. Other eclogite samples are less retrogressed and are composed of garnet, omphacite, diopside and rutile, and minor hypersthene, hornblende, plagioclase, quartz and ilmenite. Despite the strongly pervasive granulite- and amphibolite-facies overprint, five metamorphic stages have been recognized for the eclogite of the NDZ (Liu et al., 2005, 2007a, 2010): (1) a granulite-facies stage, with  $P \sim 0.8$  GPa; (2) a UHP eclogite-facies stage, with  $P = 4.0$  GPa and  $T = 900\text{--}960$  °C, witnessed by the occurrence of diamond (Xu et al., 2003, 2005; Liu et al., 2007b) and oriented mineral exsolutions in garnet and clinopyroxene (Tsai et al., 2000; Xu et al., 2003, 2005; Liu et al., 2007a; Malaspina et al., 2006); (3) a HP eclogite-facies stage, with  $P = 2.0$  GPa and  $T = 800\text{--}900$  °C, characterized by the coexistence of garnet, sodic clinopyroxene or jadeite-poor omphacite and rutile with quartz; (3) a retrograde granulite-facies stage, with  $P = 1.1\text{--}1.4$  GPa and  $T = 804\text{--}857$  °C, indicated by the presence of hypersthene, plagioclase and diopside symplectite; (4) a retrograde amphibolite-facies stage, with  $P = 0.6\text{--}0.7$  GPa and  $T = 706\text{--}777$  °C.

### **3. Analytical methods**

Mineral inclusions in zircon and oriented needles in garnet and clinopyroxene were identified using Raman spectroscopy at the Continental Dynamics Laboratory,

Chinese Academy of Geological Sciences (CAGS) and confirmed using the electron microprobe analyzer (EMPA) at the Institute of Mineral Resources, CAGS in Beijing. Furthermore, minerals relevant for this study were analyzed with a JEOL JXA-8800R EMPA at the Institute of Mineral Resources, CAGS in Beijing. The analytical conditions on the Raman and EMPA were reported by Xu et al. (2005) and Liu et al. (2009). The representative mineral compositions are presented in Table 1. The representative Raman spectra of mineral inclusions in zircon are reported in Fig. 6. Mineral abbreviations in figures and tables are after Kretz (1983).

## **4. Decompression textures**

### *4.1. Double symplectites*

In the studied samples, garnets are characterized by peculiar kelyphytic rims with two distinct reaction stages here referred as “double symplectites” (Fig. 3a and b): the fine-grained inner one is composed of amphibole and plagioclase and is inferred to derive from the decomposition of garnet during the amphibolite-facies retrogression; the outer symplectite between garnet and clinopyroxene or omphacite is mainly composed of very fine-grained hypersthene, diopside and plagioclase, and formed between reacting garnet and clinopyroxene under granulite-facies conditions; this outer symplectite is restricted to a very narrow zone. Both symplectites show vermicular texture and are devoid of mica, suggesting that the retrogressive reaction or decompression breakdown developed under fast exhumation rates and anhydrous conditions, such that the neoblastic minerals had insufficient time for recrystallization

and occurred as symplectite, *i.e.* a rapid retrograde metamorphic process from eclogite- via granulite- and amphibolite-facies conditions (Liu et al., 2005, 2007a).

#### 4.2. Needle exsolutions

A most spectacular feature of the eclogite in the NDZ is the ubiquitous occurrence of oriented needle exsolutions of different phases in clinopyroxene, garnet and apatite (Figs. 4a-d and 5a).

Oriented rods or needles in clinopyroxene can be divided into three types depending on their mineral assemblage. Type 1 clinopyroxene contains quartz rods (Fig. 4a), which have been considered to be the evidence for the prior existence of a supersilicic omphacite stable at UHP conditions ( $\geq 2.5$  GPa) (cf. Liou et al., 1998 for a review), and occur as discrete grains in the matrix or as inclusions in garnet. Type 2 clinopyroxene contains quartz + sodic plagioclase + orthopyroxene + amphibole needles (Fig. 4b). Type 3 clinopyroxene contains quartz + orthopyroxene lamellae (Fig. 4c). All these microtextures show that the precursor clinopyroxene was rich in Si and Na during the peak metamorphic conditions. Worth of note is that ultrahigh-temperature (UHT) conditions ( $> 900$  °C; Harley, 1998) may be suggested by orthopyroxene lamellae from clinopyroxene (type 3) as reported by Nakano et al. (2007). However, although UHT metamorphic conditions of 905–917 °C can be inferred by applying Cpx-Opx geothermometry (Wood and Banno, 1973; Wells, 1977) to the hypersthene lamellae and the host clinopyroxene at their contact, more data are needed to better constrain this event and to define its geological significance.

Most of oriented rods or needles in garnet are rutile; few garnet crystals host rutile + clinopyroxene + amphibole + apatite needles (Fig. 4d), which attest to the presence of a Si-Ti-Na-P-rich precursor garnet phase (majorite) stabilized at UHP conditions ( $> 5\text{--}7$  GPa) (Ye et al., 2000a; Mposkos and Kostopoulos, 2001; Song et al., 2005).

Needle exsolutions in apatite are made of pyrrhotite. The studied samples contain two types of apatite: type 1 is fluor (F)-apatite and occurs as inclusion in garnet or in the matrix, and is rich in F ( $\sim 3$  wt %) and devoid of any exsolutions; type 2 apatite is relatively poor in F ( $< 1$  wt %) and is typically associated with the oriented pyrrhotite exsolution (Figs. 3d and 5a; Table 1). Type 1 apatite make a stable assemblage with the peak minerals such as garnet, omphacite, rutile and coesite (see below) and hence formed at UHP metamorphic conditions. The UHP origin of type 1 apatite is also supported by its high F contents because it has been demonstrated that F in apatite increases with metamorphic grade and pressure (Spear and Pyle, 2002). In contrast, type 2 apatite may be the result of breakdown during decompression.

#### *4.3. Quartz inclusions*

Quartz inclusions in garnet from the studied eclogites always show sub-rounded to elliptical shapes and occur both as single crystals, and as polycrystalline aggregates 50 to 150  $\mu\text{m}$  in size (Figs. 3c and 4f). Around single large monocrystalline and polycrystalline quartz inclusions ( $> 30$   $\mu\text{m}$ ), tensional cracks are typically developed radiating into garnet (Fig. 3c), whereas only a few irregular joints are present around



very small ( $< 10\ \mu\text{m}$ ) quartz inclusions. For other mineral inclusions, radial fracturing of garnet is much less common. The observed textural characters of the quartz inclusions are commonly indicated as decompression features and are often considered as diagnostic for the identification of quartz pseudomorphs after coesite within a rigid host (Chopin, 1984).

#### 4.4. *Coesite relics*

Zircon is probably the best container for relict UHP metamorphic phases because of its chemical inactivity and extreme stability over a wide P–T interval (e.g., Chopin and Sobolev, 1995; Liou et al., 1998; Tabata et al., 1998; Parkinson and Katayama, 1999). It has been widely used to determine the presence of UHP phases, including coesite and diamond in country rock gneisses and to establish the P–T path of deeply subducted terranes (e.g., Tabata et al., 1998; Katayama et al., 2000b; Ye et al., 2000b; Liu et al., 2001, 2002, 2007b).

Recently Liu et al. (2010) analyzed by Raman spectroscopy, a number of quartz inclusions in zircon from one of the retrograded eclogite studied in this paper (sample 03LT1-1) and found a strong peak of quartz at  $466\ \text{cm}^{-1}$  but also a weak peak of coesite at  $521\ \text{cm}^{-1}$  (Fig. 6). Such peaks represent fundamental vibrations of coesite along with the typical quartz vibration as reported from quartz-transformed coesite in UHP rocks (Ghiribelli et al., 2002; Liu et al., 2002; Zhang et al., 2005). This spectral feature strongly support the presence of trace relic coesite in quartz, which might have escaped its complete transformation during decompression.

## 5. Discussion and interpretation

### 5.1. Coesite and other indicators of UHP metamorphism

The discovery of coesite in crustal rocks (Chopin, 1984; Smith, 1984) first introduced a very powerful indicator of UHP metamorphism. Unlike diamonds, the presence of only one coesite inclusion undoubtedly provides evidence for an UHP origin of the host mineral (Korsakov et al., 2009). However, due to the reaction kinetics of the coesite to quartz transformation (Mosenfelder and Bohlen, 1997; Perrillat et al., 2003 and references therein), fresh coesite rarely survives even when it is included in robust minerals. In most cases, relics of coesite are present in the core of polycrystalline quartz aggregates, surrounded by a radial crack pattern (e.g., Mosenfelder et al., 2005). These partial or complete pseudomorphs of quartz after coesite are ubiquitous over a wide range of lithologies in various orogens of different ages, and are typically involved in a multistage metamorphic history related to exhumation. These pseudomorphs are also typically used together with other mineralogical indicators to prove that the rocks experienced UHP metamorphic conditions.

In contrast to other sectors of the NDZ, micro-diamonds have not been found in the Luotian dome in the southwestern part of the NDZ; however, there are other clues supporting the UHP metamorphism over the whole unit.

Firstly, the occurrence of mineral quartz exsolutions. Although Page et al. (2005) suggested that the presence of quartz rods in clinopyroxene does not require UHP

313 metamorphism, relic coesite rods in eclogites of Tianshan in China (Zhang et al. 2005)  
314 clearly demonstrated that SiO<sub>2</sub> exsolution in clinopyroxene occurred in the coesite  
315 stability field and that most exsolved coesite rods in clinopyroxene transformed to  
316 quartz during retrograde metamorphism with only minor amounts of remnant coesite  
317 being present. Dobrzhinetskaya et al. (2002) proposed that oriented SiO<sub>2</sub> precipitation  
318 in omphacite from eclogite in the Alpe Arami garnet peridotite massif, Switzerland,  
319 occurred at P-T conditions of 7.0 GPa and 1100 °C. Experimental studies also  
320 indicated that supersilicic clinopyroxene formed under HP-UHP and HT conditions of  
321 2.5–3.2 GPa at 1400–1500 °C (Wood and Henderson 1978), 3.5–7.0 GPa at 1200 °C  
322 (Zharikov et al. 1984), and 2–3 GPa at 1200–1400 °C (Gasparik 1986). Mao (1971)  
323 found that at 4 GPa and 1100–1700 °C, clinopyroxene contains 7.5 wt% excess SiO<sub>2</sub>,  
324 and its excess SiO<sub>2</sub> increases with pressure. Therefore, based on natural and  
325 experimental data, quartz exsolution in clinopyroxene are fully compatible with, and  
326 strongly suggest, UHP and HT conditions because natural rocks containing  
327 clinopyroxene with quartz rods usually coexist with garnet or zircon containing  
328 coesite, or its quartz pseudomorph, or micro-diamond inclusions (e.g., Xu et al., 1992,  
329 2005; Zhang et al., 1995; Liou et al., 1998; Katayama et al. 2000; Liu et al., 2007b).  
330 In this regard, oriented rods or needles in clinopyroxene from the eclogites in the  
331 Luotian Dome most likely witness a former UHP stage as suggested by Tsai and Liou  
332 (2000), and this microtextural evidence is fully consistent with the presence of relic  
333 coesite in zircon from one of the studied samples. Additionally, garnet from the  
334 studied eclogites contains rutile + clinopyroxene + amphibole + apatite oriented

needles, which were ascribed by a number of authors (Ye et al., 2000a; Mposkos and Kostopoulos, 2001; Song et al., 2005) to be precursors of UHP condition at  $P > 5\text{--}7$  GPa.

Another important evidence for UHP metamorphism in the studied area is the occurrence of quartz and relic coesite inclusions. Among all studied quartz inclusions in garnet, the majority is of the monocrystalline type (Fig. 3c), some are of the polycrystalline type (2 to 5 grains per inclusion, Fig. 4e and f), and only one inclusion within a zircon of sample 03LT1-1 gives Raman evidence of relic coesite based on the presence of an additional band at  $521\text{ cm}^{-1}$  in the spectrum (Fig. 6c), corresponding to the most intense fundamental vibration in coesite. Numerous studies on natural coesite samples have documented that fundamental coesite and quartz frequencies can be present in single coesite spectra (Boyer et al., 1985; Ghiribelli et al., 2002; Liu et al., 2002; Zhang et al., 2005). This apparent discrepancy can be easily explained when the coesite grains are smaller than the laser beam to include some of the surrounding quartz, or because of incipient coesite transformation to quartz, although in some cases this effect can be induced by the laser itself. Moreover, a fracture goes through the coesite/quartz inclusions in zircon from sample 03LT1-1 (Fig. 6a), most likely resulting in the transformation of coesite to quartz.

In addition, a number of monocrystalline and polycrystalline quartz inclusions are enclosed in garnet with well-developed radial fractures (Figs. 3c and 4e), which are indicated as decompression features and are often considered to be diagnostic for the identification of quartz pseudomorphs after coesite within a rigid host (e.g., Chopin,

1984; Smith, 1984). However, although the presence of such textural features should be treated with caution because they are not unique to coesite transformation as suggested by Chopin & Sobolev (1995), the finding of relic coesite in the same rock samples supports the possibility that polycrystalline and monocrystalline quartz inclusions in garnet with radial cracks were formerly coesite, now inverted to quartz.

Therefore, the studied eclogites show radial fractures within garnet around quartz pseudomorphs and rare coesite relics documenting that they underwent UHP metamorphism, whereas multiple-phase needles in garnet show that the eclogites suffered UHP metamorphism with a possible pressure > 5–7 GPa. Other mineralogical and microtextural indicators, such as exsolved needles in clinopyroxene and F-apatite, are also present in the eclogite from the studied area and support peak UHP conditions.

## *5.2. Factors helping the poor preservation of UHP traces*

Rapid exhumation for most UHP terranes has now been confirmed with geochronological data (e.g., Gebauer et al. 1997; Rubatto and Hermann 2001; Carswell et al. 2003; Mosenfelder et al., 2005 and references therein) and stable isotope data (Zheng et al., 2003). This geodynamic mechanism also help to some extent the preservation of coesite and related UHP phases. However, according to many authors (e.g., Liou and Zhang, 1996; Liou et al., 1997; Mosenfelder et al. 2005 and references therein), the preservation of coesite and other UHP evidences may depend on many factors including the rigidity of the host mineral, the P-T conditions

and path of metamorphic crystallization, the rate of exhumation, continuous cooling during decompression, and prevention of fluid infiltration into the host mineral until fracturing at low temperatures or the presence of fluids during retrogression. Perhaps the most vital factor for the survival of coesite is its inclusion in strong host phases, such as garnet and zircon, which can act as “pressure vessels” and sustain an overpressure on the inclusion, inhibiting the volume increase necessary to transform it to quartz. That is to say, mineral inclusions in zircon or garnet may witness peak metamorphism or earlier stages of metamorphism (Chopin, 2003).

The petrologic observations described in this paper clearly reveal that the studied eclogites suffered a complex metamorphic evolution resulting in multistage decompression related to exhumation, which most likely obliterated almost all of the earlier evidence for coesite and UHP metamorphism. As a consequence, so far few evidence for UHP metamorphism has been found in the NDZ. The poor preservation of UHP conditions in the eclogites and orthogneisses of the NDZ might be also favored by the presence of partial melts during decompression from coesite- or diamond- to granulite-facies conditions as experimentally documented by Hermann et al. (2001), which is supported by the evidence for partial melting of the eclogites (Liu Y.-C. et al., unpublished data). Unfortunately, the textures of the orthogneisses cannot be used to prove this hypothesis because of their felsic composition and the subsequent, pervasive granulite- and amphibolite-facies overprint.

The P-T path followed by the eclogites of the NDZ (Fig. 7) maintained high- to very-high-temperature conditions for a long way from the early stages of uplift at

UHP conditions to HP granulite-facies overprinting. This peculiar P-T evolution, together with the influx of later abundant fluid circulation during the amphibolite-facies retrogression, may contribute to explain the extremely rare preservation of coesite and related UHP relics (Liou and Zhang, 1996; Mosenfelder et al., 2005). The early post-peak P-T history of the NDZ eclogites (Liu et al., 2000, 2007a; Xu et al., 2000), allowed to retain equilibration temperatures exceeding 850 °C during exhumation up to lower crustal levels (1.1-1.4 GPa and 804–857 °C); the following evolution is marked by both temperature and pressure decrease (Fig. 7). In such a P-T path, characterized mainly by isothermal decompression, the difference between internal and external pressure of garnet reaches its maximum at the end of isothermal decompression, and garnet fracturing would occur at high temperatures (> 800°C), leading to rapid diffusion (Nakano et al., 2007) and slow cooling with a complete breakdown of coesite (Ghiribelli et al., 2002; Faryad et al., 2010) or even the possibility that in most cases coesite would have been consumed (Tsai and Liou, 2000) at such HT condition during the granulite-facies overprint.

### *5.3. Implications for the exhumation of UHP rocks*

The data presented in this paper show that the eclogites from the Luotian dome in the NDZ experienced a multistage and relatively slow cooling process at high-T conditions after peak UHP metamorphism, with a widespread HP granulite-facies overprinting. In contrast, the CDZ and SDZ underwent rapid exhumation and cooling from the eclogite-facies to the amphibolite-facies metamorphic stage without any

granulite-facies overprinting. As a result, the eclogites in the NDZ are characterized by peculiar multiple decompression microtextures, and do not preserve almost any evidence of UHP metamorphism because of complete retrogression of UHP minerals due to action of fluids during the late amphibolite-facies overprinting and/or to the complete decomposition of coesite and related minerals during high-T exhumation. Therefore, the Dabie orogen comprises three eclogite-bearing terranes (i.e., the NDZ, CDZ and SDZ), all subjected to UHP metamorphism, characterized by different evolutionary processes and exhumation histories. This conclusion further supports the model for multi-slice and multistage successive exhumation of UHP metamorphic rocks proposed by Liu et al. (2007b).

## 6. Conclusions

The eclogites from the NDZ underwent a prolonged exhumation history from eclogite- via granulite- and amphibolite-facies conditions, resulting in the common formation of peculiar multistage decompression textures and the extremely rare preservation of former UHP metamorphic evidence. The studied eclogites were strongly affected by multiple recrystallization processes during exhumation, thus making the determination of peak metamorphic conditions particularly difficult. However, the occurrence of radial cracks around quartz inclusions and of polycrystalline quartz inclusions in garnet, most likely suggests that these inclusions formerly were coesite, now inverted to quartz. This hypothesis is also supported by the presence of relic coesite which has been confirmed by Raman spectroscopy.



Moreover, the occurrence of rutile + clinopyroxene + amphibole + apatite oriented needles in garnet open the possibility of peak pressure exceeding 5-7 GPa in the NDZ. An important consequence of this study is that if the eclogites from the Luotian dome suffered UHP metamorphism, they are perfectly comparable to those from the northeastern part of the NDZ; in this respect, the NDZ wholly experienced Triassic UHP metamorphism as a coherent unit.

## **Acknowledgements**

This research was financially supported by the National Basic Research Program of China (2009CB825002), the National Natural Science Foundation of China (40921002, 40973043 and 40572035), the PhD Foundation of the Ministry of Education of China (200803580001) and the Chinese Academy of Sciences (kzcx2-yw-131). We would like to thank Prof. Shuguang Li for his valuable discussion and helpful suggestions on this study. The authors are also grateful to Ms. Ling Yan for the Raman analysis.

## References

- Ayers, J.C., Dunkle, S., Gao, S., Miller, C.F., 2002. Constraints on timing of peak and retrograde metamorphism in the Dabie Shan ultrahigh-pressure metamorphic belt, east-central China, using U-Th-Pb dating of zircon and monazite. *Chemical Geology* 186, 315–331.
- Boyer, H., Smith, D.C., Chopin, C., Lasnier, B., 1985. Raman microprobe (RMP) determinations of natural and synthetic coesite. *Physics and Chemistry of Minerals* 12, 45–48.
- Bryant, J.L., Ayers, J.C., Gao, S., Miller, C.F., Zhang, H., 2004. Geochemical, age, and isotopic constraints on the location of the Sino-Korean/Yangtze suture and evolution of the Northern Dabie Complex, east central China. *Geological Society of America Bulletin* 116, 698–717.
- Carswell, D.A., Brueckner, H.K., Cuthbert, S.J., Mehta, K., O'Brien, P.J., 2003. The timing of stabilisation and the exhumation rate for ultra-high pressure rocks in the Western Gneiss Region of Norway. *Journal of Metamorphic Geology* 21, 601–612.
- Chavagnac, V., Jahn, B.M., 1996. Coesite-bearing eclogites from the Bixiling complex. Dabie Mountains, China: Sm–Nd ages, geochemical characteristics and tectonic implications. *Chemical Geology* 133, 29–51.
- Chopin, C., 1984. Coesite and pure pyrope in high-grade blue schists of the western Alps: a first record and some consequences. *Contributions to Mineralogy and Petrology* 86, 107–118.

483 Chopin, C., Sobolev, N.V., 1995. Principal mineralogical indicators of UHP in crustal  
 484 rocks, in: Coleman R.G. and Wang X. (eds), Ultrahigh pressure metamorphism,  
 485 Cambridge University Press, pp. 96–133.

486 Chopin, C., 2003. Ultrahigh-pressure metamorphism: tracing continental crust into the  
 487 mantle. *Earth and Planetary Science Letters* 212, 1–14.

488 Dobrzhinetskaya, L., Schweinehage, R., Massonne, H.J., Green, H.W., 2002. Silica  
 489 precipitates in omphacite from eclogite at Alpe Arami, Switzerland: evidence of  
 490 deep subduction. *Journal of Metamorphic Geology* 20, 481–492.

491 Faryad, S.W., 2009. The Kutná Hora Complex (Moldanubian zone, Bohemian Massif):  
 492 A composite of crustal and mantle rocks subducted to HP/UHP conditions. *Lithos*  
 493 109, 193–208.

494 Faryad, S.W., Nahodilová, R., Dolejš, D., 2010. Incipient eclogite facies  
 495 metamorphism in the Moldanubian granulites revealed by mineral inclusions in  
 496 garnet. *Lithos* 114, 54–69.

497 Gasparik, T., 1986. Experimental study of subsolidus phase relations and mixing  
 498 properties of clinopyroxene in silica-saturated system CaO-MgO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>.  
 499 *American Mineralogist* 71, 686–693.

500 Gayk, T., Kleinschrodt, R., Langosch, A., Seidel, E., 1995. Quartz exsolution in  
 501 clinopyroxene of high-pressure granulite from the Munchberg Massif. *European*  
 502 *Journal of Mineralogy* 7, 1217–1220.

503 Gebauer, D., Schertl, H.-P., Brix, M., Schreyer, W., 1997. 35 Ma old  
 504 ultrahigh-pressure metamorphism and evidence for very rapid exhumation in the

505 Dora Maira massif, Western Alps. *Lithos* 41, 5–24.

506 Ghiribelli, B., Frezzotti, M.-L., and Palmeri, R., 2002. Coesite in eclogites of the  
 507 Lanterman Range (Antarctica): Evidence from textural and Raman studies.  
 508 *European Journal of Mineralogy* 14, 355–360.

509 Hacker, B.R., Ratschbacher, L., Webb, L.E., Ireland, T.R., Walker, D., Dong, S., 1998.  
 510 U/Pb zircon ages constrain the architecture of the ultrahigh-pressure  
 511 Qinling-Dabie orogen, China. *Earth and Planetary Science Letters* 161, 215–230.

512 Harley, S.L., 1998. On the occurrence and characterization of ultrahigh-temperature  
 513 (UHT) crustal metamorphism. In: *What Controls Metamorphism and*  
 514 *Metamorphic Reactions?* In: Treloar, P.J., O'Brien, P. (eds.), Special Publication  
 515 *Geological Society of London* 138, 75–101.

516 Hermann, J., Rubatto, D., Korsakov, A., and Shatsky, V.S., 2001. Multiple zircon  
 517 growth during fast exhumation of diamondiferous, deeply subducted continental  
 518 crust (Kokchetav massif, Kazakhstan). *Contributions to Mineralogy and*  
 519 *Petrology* 141, 66–82.

520 Katayama, I., Parkinson, C.D., Okamoto, K., Nakajima, Y., Maruyama, S., 2000a.  
 521 Supersilicic clinopyroxene and silica exsolution in UHPM eclogite and pelitic  
 522 gneiss from the Kokchetav massif, Kazakhstan. *American Mineralogist*, 85,  
 523 1368–1374.

524 Katayama, I., Zayachkovsky, A.A., Maruyama, S., 2000b. Prograde  
 525 pressure-temperature records from inclusions in zircons from ultrahigh-pressure –  
 526 high-temperature rocks of the Kokchetav Massif, northern Kazakhstan. *Island Arc*

527 9, 417–427.

528 Korsakov, A.V., Perraki, M., Zhukov, V.P., Gussem, K.D., Vandenabeele, P.,  
529 Tomilenko, A.A., 2009. Is quartz a potential indicator of ultrahigh-pressure  
530 metamorphism? Laser Raman spectroscopy of quartz inclusions in  
531 ultrahigh-pressure garnets. *European Journal of Mineralogy* 21, 1313–1323.

532 Kretz, R., 1983. Symbols for rock-forming mineral. *American Mineralogist* 68,  
533 277–279.

534 Li, S., Xiao, Y., Liu, D., Chen, Y., Ge, N., Zhang, Z., Sun, S. S., Cong, B., Zhang, R.,  
535 Hart, S.R., Wang, S., 1993. Collision of the North China and Yangtze blocks and  
536 formation of coesite-bearing eclogites: timing and processes. *Chemical Geology*  
537 109, 89–111.

538 Li S., Jagoutz E., Chen Y., Li Q., 2000. Sm-Nd and Rb-Sr isotope chronology of  
539 ultrahigh-pressure metamorphic rocks and their country rocks at Shuanghe in the  
540 Dabie Mountains, central China. *Geochimica Cosmochimica Acta* 64, 1077–1093.

541 Li, S., Huang, F., Zhou, H., Li, H., 2003. U-Pb isotopic compositions of the ultrahigh  
542 pressure metamorphic (UHPM) rocks from Shuanghe and gneisses from Northern  
543 Dabie zone in the Dabie Mountains, central China: Constraint on the exhumation  
544 mechanism of UHPM rocks. *Science in China (Series D)* 46, 200–209.

545 Li, X.-P., Zheng, Y.-F., Wu, Y.-B., Chen, F.K., Gong, B., Li, Y.-L., 2004. Low-T  
546 eclogite in the Dabie terrane of China: petrological and isotopic constrains on  
547 fluid activity and radiometric dating. *Contributions to Mineralogy and Petrology*  
548 148, 443–470.

- 549 Liou, J.G., Zhang, R.Y., 1996. Occurrences of intergranular coesite in ultrahigh-P  
550 rocks from the Sulu region, eastern China: implications for lack of fluid during  
551 exhumation. *American Mineralogist* 81, 1217–1221.
- 552 Liou, J.G., Zhang, R.Y., Jahn, B.M., 1997. Petrology, geochemistry and isotope data  
553 on a ultrahigh-pressure jadeite quartzite from Shuanghe, Dabie Mountains,  
554 east-central China. *Lithos* 41, 50–78.
- 555 Liou, J. G., Zhang, R. Y., Ernst, W. G., Rumble, D., Maruyama, S., 1998.  
556 High-pressure minerals from deeply subducted metamorphic rocks. Review of  
557 Mineralogy 37, 33–96.
- 558 Liu, F.L., Xu, Z.Q., Liou, J.G., Katayama, I., Masago, H., Maruyama, S., Yang, J.,  
559 2002. Ultrahigh-pressure mineral inclusions in zircons from gneissic core samples  
560 of the Chinese Continental Scientific Drilling Site in eastern China. *European*  
561 *Journal of Mineralogy* 14, 499–512.
- 562 Liu, J.B., Ye, K., Maruyama, S., Cong, B.L., Fa, H.R., 2001. Mineral inclusions in  
563 zircon from gneisses in the ultrahigh-pressure zone of the Dabie Mountains,  
564 China. *The Journal of Geology* 109, 523–535.
- 565 Liu, Y.-C., Li, S., Xu, S., Li, H., Jiang, L., Chen, G., Wu, W., Su, W., 2000. U-Pb  
566 zircon ages of the eclogite and tonalitic gneiss from the northern Dabie  
567 Mountains, China and multi-overgrowths of metamorphic zircons. *Geological*  
568 *Journal of China Universities* 6, 417–423 (in Chinese with English abstract).
- 569 Liu, Y.-C., Li, S., Xu, S., Jahn, B.-M., Zheng, Y., Zhang, Z., Jiang, L., Chen, G., Wu,  
570 W., 2005. Geochemistry and geochronology of eclogites from the northern Dabie

571 Mountains, central China. *Journal of Asian Earth Sciences* 25, 431–443.

572 Liu, Y.-C., Li, S., Gu, X., Xu, S., Chen, G., 2007a. Ultrahigh-pressure eclogite  
 573 transformed from mafic granulite in the Dabie orogen, east-central China. *Journal*  
 574 *of Metamorphic Geology* 25, 975–989.

575 Liu, Y.-C., Li, S., Xu, S., 2007b. Zircon SHRIMP U-Pb dating for gneiss in northern  
 576 Dabie high T/P metamorphic zone, central China: Implication for decoupling  
 577 within subducted continental crust. *Lithos* 96, 170–185.

578 Liu, Y.-C., Li, S., 2008. Detachment within subducted continental crust and  
 579 multi-slice successive exhumation of ultrahigh-pressure metamorphic rocks:  
 580 Evidence from the Dabie-Sulu orogenic belt. *Chinese Science Bulletin* 53,  
 581 3105–3119.

582 Liu, Y.-C., Wang, A., Rolfo, F., Groppo, C., Gu, X., Song, B., 2009. Geochronological  
 583 and petrological constraints on Palaeoproterozoic granulite facies metamorphism  
 584 in southeastern margin of the North China Craton. *Journal of Metamorphic*  
 585 *Geology* 27, 125–138.

586 Liu, Y.-C., Gu, X., Li, S., Hou, Z., Song, B., 2010. Multistage metamorphic events in  
 587 eclogites from the North Dabie high-T/UHP Complex zone, central China:  
 588 evidence from zircon U-Pb age, trace element and mineral inclusion. Submitted.

589 Malaspina, N., Hermann, J., Scambelluri, M., Compagnoni, R., 2006. Multistage  
 590 metasomatism in ultrahigh-pressure mafic rocks from the North Dabie Complex  
 591 (China). *Lithos* 90, 19–42.

592 Mao, H.K., 1971. The system jadeite (NaAlSi<sub>2</sub>O<sub>6</sub>)-anorthite (CaAl<sub>2</sub>Si<sub>2</sub>O<sub>8</sub>) at high

593 pressure. Carnegie Institute Year Book 69, 163–168.

594 Mosenfelder, J.L., Bohlen, S.R., 1997. Kinetics of the coesite to quartz transformation.

595 Earth and Planetary Science Letters 153, 133–147.

596 Mosenfelder, J.L., Schertl, H.-P., Smyth, J.R., Liou, J.G., 2005. Factors in the

597 preservation of coesite: The importance of fluid infiltration. American

598 Mineralogist 90, 779–789.

599 Mposkos, E.D., Kostopoulos, D.K., 2001. Diamond, former coesite and supersilicic

600 garnet in metasedimentary rocks from the Greek Rhodope: a new

601 ultrahigh-pressure metamorphic province established. Earth and Planetary

602 Science Letters 192, 497–506.

603 Nakano, N., Osanai, Y., Owada, M., 2007. Multiple breakdown and chemical

604 equilibrium of silicic clinopyroxene under extreme metamorphic conditions in the

605 Kontum Massif, central Vietnam. American Mineralogist 92, 1844–1855.

606 Okay, A.I., Xu S., Sengör, A.M.C., 1989. Coesite from the Dabie Shan eclogites,

607 central China. European Journal of Mineralogy 1, 595–598.

608 Okay, A.I., 1993. Petrology of a diamond and coesite-bearing metamorphic terrain:

609 Dabie Shan, China. European Journal of Mineralogy 5, 659–675.

610 Okay, A.I., Sengör, A.M.C., Satir, M., 1993. Tectonics of an ultrahigh-pressure

611 metamorphic terrane: the Dabie Shan/Tongbai orogen, China. Tectonics 12,

612 1320–1334.

613 Page, F.Z., Essene, E.J., Mukasa, S.B., 2005. Quartz exsolution in clinopyroxene is

614 not proof of ultrahigh pressures: Evidence from eclogites from the Eastern Blue



615 Ridge, Southern Appalachians, U.S.A. *American Mineralogist* 90, 1092–1099.

616 Parkinson, C. D., Katayama, I., 1999. Present day ultrahigh-pressure conditions of  
 617 coesite inclusions in zircon and garnet: evidence from laser Raman  
 618 microspectroscopy. *Geology* 27, 979–982.

619 Perrillat, J.P., Daniel, I., Lardeaux, J.M., Cardon, H., 2003. Kinetics of the  
 620 coesite–quartz transition: application to the exhumation of ultrahigh-pressure  
 621 rocks. *Journal of Petrology*, 44, 773–788.

622 Rolfo, F., Compagnoni, R., Xu, S., Jiang, L., 2000. First report of felsic whiteschist in  
 623 the ultrahigh-pressure metamorphic belt of Dabie Shan, China. *European Journal*  
 624 *of Mineralogy* 12, 883–898.

625 Rolfo, F., Compagnoni, R., Wu, W., Xu, S., 2004. A coherent lithostratigraphic unit in  
 626 the coesite-eclogite complex of Dabie Shan, China: geologic and petrologic  
 627 evidence. *Lithos* 73, 71–94.

628 Rubatto, D., Hermann, J., 2001. Exhumation as fast as subduction? *Geology* 29, 3–6.

629 Smith, D.C., 1984. Coesite in clinopyroxene in the Caledonides and its implications  
 630 for geodynamics. *Nature* 310, 641–644.

631 Song, S.G., Yang, J.S., Xu, Z.Q., Liou, J.G., Shi, R.D., 2003. Metamorphic evolution  
 632 of the coesite-bearing ultrahigh-pressure terrane in the North Qaidam, northern  
 633 Tibet, NW China. *Journal of Metamorphic Geology* 21, 631–644.

634 Song, S.G., Zhang, L., Chen, J., Liou, J.G., Niu, Y., 2005. Sodic amphibole exsolutions  
 635 in garnet from garnet-peridotite, North Qaidam UHPM belt, NW China:  
 636 Implications for ultradeep-origin and hydroxyl defects in mantle garnets.

637 American Mineralogist 90, 814–820.

638 Spear, F.S., Pyle, J.M., 2002. Apatite, Monazite, and Xenotime in metamorphic rocks.

639 Reviews in Mineralogy and Geochemistry 48, 293–335.

640 Tabata, H., Yamauchi, K., Maruyama, S., Liou J.G., 1998. Tracing the extent of a UHP

641 metamorphic terrane: Mineral-inclusion study of zircons in gneisses from the

642 Dabieshan, In: B.R. Hacker, J.G. Liou (eds.), When Continents Collide:

643 Geodynamics and Geochemistry of Ultrahigh-Pressure Rocks. Kluwer Academic

644 Publisher, pp. 261–273.

645 Tsai, C.H., Liou, J.G., 2000. Eclogite-facies relics and inferred ultrahigh-pressure

646 metamorphism in the North Dabie complex, central China. American Mineralogist

647 85, 1–8.

648 Wang, X., Liou, J.G., Mao, H.K., 1989. Coesite-bearing eclogites from the Dabie

649 Mountains in central China. Geology 17, 1085–1088.

650 Wells, R.A., 1977. Pyroxene thermometry in simple and complex systems.

651 Contributions to Mineralogy and Petrology 62, 129–139.

652 Wood, B.J., Banno, S., 1973. Garnet-orthopyroxene and orthopyroxene-clinopyroxene

653 relationship in simple and complex systems. Contributions to Mineralogy and

654 Petrology 42, 109–124.

655 Wood, B.J., Henderson, C.M.B., 1978. Compositions and unit-cell parameters of

656 synthetic non-stoichiometric tschermakitic clinopyroxenes. American

657 Mineralogist 63, 66–72.

658 Xu, S., Okay, A.I., Ji, S., Sengör, A.M.C., Su, W., Liu, Y., Jiang, L., 1992. Diamond

659 from the Dabie Shan metamorphic rocks and its implication for tectonic setting.  
 660 Science 256, 80–82.

661 Xu, S., Liu, Y.-C., Su, W., Wang, R., Jiang, L., Wu, W., 2000. Discovery of the  
 662 eclogite and its petrography in the Northern Dabie Mountains. Chinese Science  
 663 Bulletin 45, 273–278.

664 Xu, S., Liu, Y.-C., Chen, G., Compagnoni, R., Rolfo, F., He, M., Liu, H., 2003. New  
 665 finding of micro-diamonds in eclogites from Dabie-Sulu region in central-eastern  
 666 China. Chinese Science Bulletin 48, 988–994.

667 Xu, S., Liu, Y.-C., Chen, G., Ji, S., Ni, P., Xiao, W., 2005, Microdiamonds, their  
 668 classification and tectonic implications for the host eclogites from the Dabie and  
 669 Su-Lu regions in central eastern China. Mineralogical Magazine 69, 509–520.

670 Ye K., Cong B., Ye D., 2000a. The possible subduction of continental material to  
 671 depths greater than 200 km. Nature 407, 734–736.

672 Ye, K., Yao, Y., Katayama, I., Cong, B., Wang, Q., Maruyama, S., 2000b. Large areal  
 673 extent of ultrahigh-pressure metamorphism in the Sulu ultrahigh-pressure terrane  
 674 of East China: new implications from coesite and omphacite inclusions in zircon  
 675 of granitic gneiss. Lithos 52, 157–164.

676 Zhang, H., Gao, S., Zhong, Z., Zhang, B., Zhang, L., Hu, S., 2002a. Geochemical and  
 677 Sr–Nd–Pb isotopic compositions of Cretaceous granitoids: constraints on tectonic  
 678 framework and crustal structure of the Dabieshan ultrahigh-pressure metamorphic  
 679 belt, China. Chemical Geology 186, 281–299.

680 Zhang L., Ellis D.J., Jiang W., 2002b. Ultrahigh pressure metamorphism in western

Tianshan, China, part I: evidences from the inclusion of coesite pseudomorphs in garnet and quartz exsolution lamellae in omphacite in eclogites. *American Mineralogist* 87, 853–860.

Zhang, L.F., Song, S., Liou, J.G., Ai, Y., Li, X., 2005. Relict coesite exsolution in omphacite from Western Tianshan eclogites, China: *American Mineralogist* 90, 181–186.

Zhang, R.Y., Hirajima, T., Banno, S., Cong, B., Liou, J.G., 1995. Petrology of ultrahigh pressure rocks from the southern Sulu region, eastern China. *Journal of Metamorphic Geology* 13, 659–675.

Zhang, R.Y., Zhai, S.M., Fei, Y.W., Liou, J.G., 2003. Titanium solubility in coexisting and clinopyroxene at very high pressure: the significance of exsolved rutile in garnet. *Earth and Planetary Science Letters* 216, 591–601.

Zhang, R.Y., Liou, J.G., Ernst, W.G., 2009. The Dabie–Sulu continental collision zone: A comprehensive review. *Gondwana Research* 16, 1–26.

Zhao, Z., Zheng, Y., Wei, C., Chen, F., Liu, X., Wu, F., 2008. Zircon U–Pb ages, Hf and O isotopes constrain the crustal architecture of the ultrahigh-pressure Dabie orogen in China. *Chemical Geology* 253, 222–242.

Zharikov, V.A., Ishbulatov, R.A., Chudinovskikh, L.T., 1984. High-pressure clinopyroxenes and eclogite barrier. *Soviet Geology and Geophysics* 25, 53–61.

Zheng, Y., Fu, B., Gong, B., Li, L., 2003. Stable isotope geochemistry of ultrahigh pressure metamorphic rocks from the Dabie–Sulu orogen in China: implications for geodynamics and fluid regime. *Earth Science Reviews* 62, 105–161.

## Figure captions

**Figure 1** Schematic geological map of the Dabie orogen. Sample localities with sample numbers are described in detail in the text. BZ = Beihuaiyang zone, NDZ = North Dabie complex zone, CDZ = Central Dabie UHP metamorphic zone, SDZ = South Dabie low-T eclogite zone, SZ = Susong complex zone, HMZ = Huwan mélangé zone, HZ = Hong'an low-T eclogite zone, DC = amphibolite-facies Dabie complex, XMF= Xiaotian-Mozitan fault, WSF= Wuhe-Shuihou fault, HMF = Hualiangting-Mituo fault, TSF= Taihu-Shanlong fault, TLF = Tan-Lu fault. The inset shows the location of Fig. 1 within the Triassic Qinling—Dabie—Su-Lu collision orogen in central China.

**Figure 2** Field occurrence of eclogite band (a) and lens (b) from the Luotian dome in the southwestern part of the North Dabie complex zone.

**Figure 3** Photomicrographs of eclogite from the Luotian dome in the Dabie orogen. a. Omphacite inclusion in garnet with two generations of symplectites (Hy+Di+Pl and Hbl+Pl+Mt), sample 06LT4-2; b. Omphacite, rutile and quartz inclusions in garnet, rimmed by distinctive double symplectites, sample 07LT6-1; c. quartz inclusion in garnet with well-developed radial fractures, sample 07LT6-1; d. Apatite (Ap) occurs as inclusion in garnet and in the matrix, the former is fluor-apatite (F-Ap) and the latter contains oriented pyrrhotite exsolutions (Po), sample 06LT4-2.

**Figure 4** Photomicrographs (a and d–f) and Back scattered electron (BSE) images (b and c) of eclogite from the Luotian dome showing multiple decompression textures. a. Diopside inclusion with quartz rods in garnet, sample 03LT1-1; b. Qtz + Pl + Hbl + Hy oriented needles in clinopyroxene, sample 03LT8-1; c. Qtz + Hy oriented needles in clinopyroxene, sample 06LT3-2; d. Cpx + Rt + Hbl + Ap oriented needles in garnet, sample 03LT1-1; e. Quartz after coesite in garnet, sample 09LT2-1; f. Cross-polarized light image of (e), showing polycrystalline quartz inclusions in garnet.

**Figure 5** a. Photomicrograph showing needle pyrrhotite (Po) exsolutions in apatite (Ap) and Cpx with quartz needles, sample 06LT3-2; b. The EDS spectra of exsolution rods in apatite from (a).


**Figure 6** Photomicrographs (a and b) of coesite (Cs) and quartz (Qtz) inclusion in zircon (Zr) and their mixed Raman spectra (c); zircon separated from the eclogite sample 03LT1-1 from the Luotian dome. The black circle in (a) is LA-ICPMS dating analysis with available  $^{206}\text{Pb}/^{208}\text{U}$  age shown.

**Figure 7** Multistage exhumation P-T trajectory of the NDZ eclogites, based on available data from the north-eastern portion of the NDZ (Malapina et al., 2006) and from the Luotian dome (Liu et al., 2007a).

747    **Table captions**

748    Table 1 Electron microprobe analyses of representative minerals from the eclogites in  
749    the Luotian dome (wt%)

**Figure 1**  
[Click here to download high resolution image](#)



**Figure 1**



**Figure 2**  
[Click here to download high resolution image](#)



**Figure 2**



Figure 3  
[Click here to download high resolution image](#)




Figure 3

Figure 4  
[Click here to download high resolution image](#)




Figure 4



Figure 5  
[Click here to download high resolution image](#)





Figure6  
[Click here to download high resolution image](#)




Figure 7  
[Click here to download high resolution image](#)




Figure 7

Table 1

Mineral	Garnet		Omphacite		Diopside	Hornblende		Plagioclase	
No.	06LT3-2	06LT4-2	06LT3-2	06LT4-2	06LT3-2	06LT3-2	06LT3-2	03LT1-1	06LT3-2
Locality	m	m	i	i	ng	nc	ng	nc	nc
<b>SiO<sub>2</sub></b>	38.88	38.43	54.45	55.28	52.7	45.12	38.65	61.2	61.46
<b>TiO<sub>2</sub></b>	0.02	0.07	0	0.04	0.1	0.08	0.73	0	0
<b>Al<sub>2</sub>O<sub>3</sub></b>	21.24	21.88	4.84	11.03	3.5	12.21	18.36	24.11	23.55
<b>FeO</b>	24.1	24.84	6.15	6.86	6.38	9.2	12.07	0	0.23
<b>Cr<sub>2</sub>O<sub>3</sub></b>	0	0	0.02	0	0.04	0.04	0	0	0
<b>MnO</b>	0.52	0.38	0.02	0.02	0.05	0.04	0.02	0.24	0.02
<b>MgO</b>	8.14	5.78	12.71	7.38	14.05	15.13	11.45	0	0.01
<b>CaO</b>	7.07	8.91	18.34	12.24	22.28	11.53	11.39	5.7	4.92
<b>Na<sub>2</sub>O</b>	0	0.12	2.95	7.33	0.83	2.27	2.89	8.71	9.05
<b>K<sub>2</sub>O</b>	0	0	0	0.01	0.01	0.03	0.02	0.01	0.02
<b>Total</b>	99.97	100.41	99.48	100.19	99.94	95.65	95.58	99.97	99.26
<b>O</b>	12	12	6	6	6	23	23	8	8
<b>Si</b>	2.985	2.964	1.986	1.967	1.938	6.526	5.71	2.723	2.749
<b>Al<sup>IV</sup></b>	0.015	0.036	0.014	0.033	0.062	1.474	2.29	1.263	1.241
<b>Al<sup>VI</sup></b>	1.905	1.951	0.194	0.430	0.09	0.606	0.904		
<b>Fe<sup>3+</sup></b>	0.102	0.09	0.028	0.106	0.025	0.626	0.787	0	0
<b>Ti</b>	0.001	0.004	0.000	0.001	0.003	0.009	0.081	0	0
<b>Fe<sup>2+</sup></b>	1.445	1.512	0.160	0.098	0.171	0.484	0.704	0.009	0.009
<b>Cr</b>	0	0	0.001	0	0.001	0.005	0	0	0
<b>Mg</b>	0.932	0.664	0.691	0.392	0.77	3.263	2.522	0	0.001
<b>Mn</b>	0.034	0.025	0.001	0.001	0.002	0.005	0.003	0	0.001
<b>Ca</b>	0.581	0.736	0.717	0.467	0.878	1.787	1.803	0.272	0.236
<b>Na</b>	0	0.018	0.209	0.506	0.059	0.637	0.828	0.751	0.785
<b>K</b>	0	0	0	0	0	0.006	0.004	0.001	0.001

Note: m, matrix; ng, oriented needle in garnet; nc, oriented needle in clinopyroxene; i, inclusion; sy, symplectite.

Continued Table 1

Mineral	Apatite		Diopside		Hypersthene		Hornblende	Plagioclase	
No.	06LT3-2	06LT4-2	07LT6-1	06LT3-2	06LT3-2	07LT6-1	07LT6-1	07LT6-1	07LT6-1
Locality	m	i	sy	m	nc	sy	sy	sy	sy
SiO <sub>2</sub>	0.00	0.00	53.28	54.39	54.89	51.22	42.26	58.82	55.19
TiO <sub>2</sub>	0.02	0.00	0.18	0.03	0.00	0.08	0.86	0.02	0.04
Al <sub>2</sub> O <sub>3</sub>	0.01	0.00	1.33	3.90	2.10	0.36	14.94	25.93	28.5
FeO	0.17	0.37	7.10	6.59	18.9	29.73	13.53	0.36	0.54
Cr <sub>2</sub> O <sub>3</sub>	0.01	0.01	0.11	0	0.01	0.01	0.16	0.00	0.00
MnO	0.02	0.00	0.10	0.03	0.13	0.47	0.06	0.02	0.01
MgO	0.08	0.04	14.81	13.99	23.42	17.75	11.15	0.02	0.02
CaO	55.69	54.45	22.61	20.35	0.50	0.59	9.33	6.98	10.15
Na <sub>2</sub> O	0.06	0.00	0.42	1.52	0.01	0.02	4.12	7.42	6.02
K <sub>2</sub> O	0.00	0.01	0.01	0.00	0.02	0.00	0.08	0.01	0.02
P <sub>2</sub> O <sub>5</sub>	41.13	41.81							
F	1.00	3.09							
Total	98.19	99.78	99.95	100.86	99.98	100.23	96.49	99.58	100.49
O			6	6	6	6	23	8	8
Si			1.969	1.974	2.018	1.968	6.200	2.635	2.480
Al <sup>IV</sup>			0.031	0.026	0.000	0.016	1.800	1.368	1.508
Al <sup>VI</sup>			0.027	0.141	0.091	0.000	0.781		
Fe <sup>3+</sup>			0.022	0.000	0.000	0.045	0.691	0.000	0.000
Ti			0.005	0.001	0.000	0.002	0.095	0.001	0.001
Fe <sup>2+</sup>			0.198	0.098	0.581	0.911	0.968	0.013	0.020
Cr			0.003	0	0.000	0.000	0.019	0.000	0.000
Mg			0.816	0.757	1.284	1.017	2.438	0.001	0.001
Mn			0.003	0.002	0.004	0.015	0.007	0.001	0.000
Ca			0.895	0.791	0.020	0.024	1.466	0.335	0.489
Na			0.030	0.107	0.001	0.001	1.172	0.645	0.524
K			0.000	0	0.001	0.000	0.015	0.001	0.001

Note: m, matrix; ng, oriented needle in garnet; nc, oriented needle in clinopyroxene; i, inclusion; sy, symplectite.